

Characters Taxon	(E) Equatorial diameter ( $\mu\text{m}$ )			(P) Polar diameter ( $\mu\text{m}$ )			P/E	Pollen shape	Mean Exoaperture length ( $\mu\text{m}$ )	Mean Exoaperture distances ( $\mu\text{m}$ )
	Min	Max	Mean	Min	Max	Mean				
<i>N. caspica</i>	19.82	32.72	26.70	28.38	43.39	34.55	1.29	subprolate	11.28	9.10
<i>N. diffusa</i>	15.20	30.00	21.56	19.87	32.00	22.66	1.06	Prolate- spheroidal	7.76	9.00
<i>N. lutea</i> var. <i>flavescens</i>	26.29	29.82	28.27	37.56	40.25	38.90	1.40	prolate	10.60	9.03
<i>N. lutea</i> var. <i>lutea</i>	26.66	30.41	29.20	35.59	42.99	39.44	1.40	prolate	10.52	8.04
<i>N. melanocar- pa</i>	24.74	28.78	26.08	30.80	34.30	32.25	1.20	subprolate	11.32	10.20
<i>N. persica</i> var. <i>persica</i>	16.52	19.34	18.44	20.04	23.51	21.98	1.19	subprolate	8.69	9.82
<i>N. persica</i> var. <i>suchtelenioides</i>	18.48	21.44	20.52	23.51	25.79	23.84	1.21	subprolate	10.20	11.18
<i>N. pulla</i>	21.38	24.36	22.65	25.92	29.42	27.50	1.21	subprolate	8.63	9.86
<i>N. rosea</i>	19.52	27.48	23.02	26.29	33.61	29.77	1.30	subprolate	11.18	10.39
<i>N. turcomanica</i>	29.85	31.49	30.35	34.35	36.39	35.41	1.16	subprolate	8.93	11.14

